

# Use of orbital cycles to improve magnetostratigraphic dating of drill core SG-1 from the Qaidam Basin (NE Tibetan Plateau) spanning nearly the last ~2.8 Ma

Christian Herb<sup>1</sup>, Andreas Koutsodendris<sup>2</sup>, Weilin Zhang<sup>1,3</sup>, Erwin Appel<sup>1</sup>, Xiaomin Fang<sup>3</sup>,  
Silke Voigt<sup>2</sup>, Jörg Pross<sup>2</sup>

<sup>1</sup> Department of Geosciences, University of Tübingen, Hölderlinstr. 12, 72074 Tübingen, Germany

<sup>2</sup> Institute of Geosciences, University of Frankfurt, Altenhöferallee 1, 60438 Frankfurt, Germany

<sup>3</sup> Institute of Tibetan Plateau Research, Chinese Academy of Sciences, P.O. Box 2871, Beijing 100085, China  
christian.herb@uni-tuebingen.de

The ~940-m-long drill core SG-1 consisting of late Pliocene-Quaternary lacustrine sediments from the western Qaidam Basin (NE Tibetan Plateau) serves as a promising archive for studying past climate change. Magnetic susceptibility ( $\chi$ ) was identified as a well-suited proxy for reconstructing past climate of the western Qaidam Basin in a high resolution (~1 ka). The causes for the  $\chi$  variation related to climate change were previously reduced to the interference of low-temperature oxidation in the catchment and a change of the sediment source regions. To bring the information of past environmental change into a meaningful time frame it is necessary to improve previous magnetostratigraphic dating (~2.8 Ma to 0.1 Ma) of drill core SG-1. Average sediment accumulation rates (SARs) estimated between geomagnetic reversals are in the order of 35 cm/ka. Time series analysis is applied for  $\chi$  to establish an orbital age model. Results of spectral analysis reveal the response of  $\chi$  to orbital forcing, including the transition from the dominating 41-ka cycle to the 100-ka cycle around the Mid-Pleistocene Transition (MPT; ~1 Ma). The observed cyclicity of  $\chi$  is used to establish a cyclostratigraphy that results in an improved age model and depth-to-age transformation of drill core SG-1. The approach includes separation of the core into intervals with different cyclic behaviour and the analysis of sliding windows.

---

Key words (for online publication): Magnetic proxies; Cyclostratigraphy; Spectral analysis; Lacustrine sediments; Paleoclimate; Qaidam Basin; Tibetan Plateau